

Bunionectomy

ORG: S-260 (ISC)

[Link to Codes](#)

MCG Health

Inpatient & Surgical Care
30th Edition

- Care Planning - Inpatient Admission and Alternatives
 - Clinical Indications for Procedure
 - Alternatives to Procedure
 - Operative Status Criteria
- Hospitalization
 - Optimal Recovery Course
 - Goal Length of Stay - **Ambulatory**
 - Extended Stay
- Discharge
 - Discharge Planning
 - Discharge Destination
- Evidence Summary
 - Background
 - Criteria
 - Alternatives
 - Length of Stay
 - Rationale
 - Related CMS Coverage Guidance
- References
- Footnotes
- Definitions
- Codes

Care Planning - Inpatient Admission and Alternatives

Clinical Indications for Procedure

- Procedure is indicated for **ALL** of the following(1)(2):**NN**
 - ☐ Symptomatic bunion (hallux valgus), as indicated by **1 or more** of the following:
 - Skin breakdown or ulceration at metatarsophalangeal joint(1)(9)
 - Difficulty walking (eg, refractory pain, imbalance, impaired stability)(3)(7)(9)
 - Inability to accommodate or modify footwear to control pain(1)(6)(10)
 - Avulsion fracture of proximal phalanx(8)
 - Malunion or nonunion of prior surgery(11)(12)(13)

- Nonoperative therapy (eg, shoe modification, bunion shield, splinting) has been tried and failed.(1)(7)(10)

Alternatives to Procedure

- Alternatives include nonoperative therapy, including:^N
 - Shoe modification(10)(14)(15)
 - Orthoses(3)(10)(15)
 - Activity adjustments(1)(10)
 - Physical therapy(3)(10)(14)
 - Toe separators(7)(16)
 - Debridement of hyperkeratotic lesions(6)
 - Oral NSAIDs(14)
 - Corticosteroid injections for local inflammation or bursitis(1)(6)
 - Splints to reduce flexible deformities(3)(10)
 - Lateral hemiepiphyodesis(9)(17)(18)
 - Fusion of first metatarsocuneiform joint(19)(20)
 - Resection arthroplasty(1)(20)

Operative Status Criteria

Goal Length of Stay: Ambulatory

Note: The definition of an ambulatory procedure depends on payer-provider contractual agreement or regulatory language (eg, CMS' Two-Midnight Rule). An ambulatory procedure may include one postoperative overnight stay in a facility; therefore, MCG's ambulatory Goal Length of Stay (GLOS) attainment calculation reports the sum of same-day and next-day postoperative discharges. Depending on various patient and procedural factors, some patients undergoing a procedure with an ambulatory GLOS require inpatient care (eg, medical necessity for hospital-based care across 2 or more postoperative midnights). Some of these factors are described in the Extended Stay section of this guideline.

- Ambulatory
- Inpatient (eg, medical necessity for hospital-based care across 2 or more postoperative midnights)
- Inpatient (Medicare patient, and specific procedure is on CMS Inpatient Only List)

Hospitalization

Optimal Recovery Course

Day	Level of Care	Clinical Status	Activity	Routes	Interventions	Medications
1	- Social Determinants of Health Assessment - OR to recovery room to discharge^[A] - Discharge planning	Hemodynamic stability Adequate supported ambulation No evidence of postoperative or surgical site infection Pain absent or managed	Ambulatory or acceptable for next level of care^[B]	- IV fluids, medications for procedure Oral hydration^[C] Oral medications or regimen acceptable for next level of care	Gait training if needed	Multimodal analgesia

- Discharge plans and education understood

- Oral diet or acceptable for next level of care

(1)(7)(9)(21)[N](#)

Recovery Milestones are indicated in **bold**.

Goal Length of Stay: Ambulatory

Note: Goal Length of Stay assumes optimal recovery, decision making, and care. Patients may be discharged to a lower level of care (either later than or sooner than the goal) when it is appropriate for their clinical status and care needs.

Extended Stay

Minimal (a few hours to 1 day), Brief (1 to 3 days), Moderate (4 to 7 days), and Prolonged (more than 7 days).

- Inpatient stay (eg, need for hospital-based care beyond postoperative day 1) may be needed for(22):
 - Failure to meet discharge criteria (recovery milestones in Optimal Recovery Course)
 - Expect brief stay extension.

See Common Complications and Conditions [ISC](#) for further information.

Discharge

Discharge Planning

- Discharge planning includes[D]:
 - Assessment of needs and planning for care, including(24)(25):
 - Develop and modify treatment plan (involving multiple providers) as needed.
 - Evaluate and address preadmission functioning as needed.
 - Evaluate and address psychosocial status issues as indicated. See Psychosocial Assessment [SR](#) for further information.
 - Evaluate and address social determinants of health (eg, housing, food). See Social Determinants of Health Screening Tool [SR](#) for further information.
 - Evaluate and address patient or caregiver preferences as indicated.
 - Identify skilled services needed at next level of care, with specific attention to(1)(26):
 - Neurovascular status assessment(27)
 - Pain management(28)
 - Wound or dressing management(29)
 - Early identification of anticipated discharge destination; options include(25)(30):
 - Home; considerations include:
 - Home safety assessment. See Home Safety Assessment [SR](#) for further information.

☐ Patient safe to go home; examples include(31)(32)(33):

- Medical status stable for patient's condition
- Functional care can safely be provided with available resources.
- Mental status stable for patient's condition
- Medication availability confirmed and reconciliation complete
- Patient/caregiver education completed with written discharge instructions provided
- Community resources identified and referrals made, as needed
- Home care arranged, if indicated
- Necessary medical equipment delivery arranged or available in home, if indicated
- Necessary medical supplies ordered, or patient/caregiver can obtain, if indicated

- Access to follow-up care

- Self-management ability if appropriate. See Activities of Daily Living (ADL) and Instrumental Activities of Daily Living (IADL) Assessment [SR](#) for further information.
- Caregiver need, ability, and availability

- Post-acute skilled care or custodial care as indicated. See Discharge Planning Tool [SR](#) for further information.

- Transitions of care plan complete, including(25)(30)(34):

- Patient and caregiver education complete.

- See Teach-Back Tool [SR](#) for further information.
- See Bunionectomy: Patient Education for Clinicians [SR](#) for further information.

☐ Medication reconciliation completion includes(35)(36):

- Compare patient's discharge list of medications (prescribed and over-the-counter) against provider's admission or transfer orders.
- Assess each medication for correlation to disease state or medical condition.
- Report medication discrepancies to prescribing provider, attending physician, and primary care provider, and ensure accurate medication order is identified.
- Provide reconciled medication list to all treating providers.
- Confirm that patient or caregiver can acquire medication.
- Educate patient and caregiver.
 - Provide complete medication list to patient and caregiver.
 - Importance of presenting personal medication list to all providers at each care transition, including all provider appointments
 - Reason, dosage, and timing of medication (eg, use "teach-back" techniques)(37)
- Encourage communication between patient, caregiver, and pharmacy for obtaining prescriptions, setting up home medication delivery, and reviewing for drug-drug interactions.
- See Medication Reconciliation Tool [SR](#) for further information.

- Plan communicated to patient, caregiver, and all members of care team, including(38):

- Inpatient care and service providers
- Primary care provider
- All post-discharge care and service providers

- Appointments planned or scheduled, which may include:

- Primary care provider(39)(40)
- Orthopedic surgeon(27)
- Podiatrist(39)

- Rehabilitation therapy services(1)(28)
- Other
- Outpatient testing and procedure plans made, which may include:
 - Other
- Referrals made for assistance or support, which may include:
 - Financial, for follow-up care, medication, and transportation(41)
 - Tobacco use treatment(42)
 - Other
- Medical equipment and supplies coordinated (ie, delivered or delivery confirmed), which may include:
 - Wound care equipment and supplies(29)(43)
 - Other

Discharge Destination

- Post-hospital levels of admission may include:
 - Home.
 - Home healthcare. See Home Care Indications for Admission Section [HC](#) in Bunionectomy guideline in Home Care.

Evidence Summary

Background

Hallux valgus, also known as a bunion, is a bony protrusion that forms on the inside of the foot near the joint of the first toe; it may benefit from surgery. Open, percutaneous, and minimally invasive surgical techniques may be used.(1)(3)(4)(5) **(EG 2)**

Criteria

The evidence for the clinical indications found in this guideline includes 10 published peer reviewed articles, 1 Cochrane systematic review, and 2 book sections.

An orthopedic surgery textbook and a narrative review support bunionectomy for patients with a bunion that causes severe symptoms (eg, pain or ulceration at the metatarsophalangeal joint, gait abnormalities, or difficulty wearing fitted shoes) that have not responded to nonoperative therapy.(1)(6) **(EG 2)** A narrative review notes that operative treatment for symptomatic bunions can be helpful to reduce deformity and the underlying deforming forces if nonoperative management has not been successful.(7) **(EG 2)**

For avulsion fracture of the proximal phalanx or grade III (complete) tear of the capsular ligamentous complex of the toe, a narrative review supports hallux valgus surgery with direct repair of the injured structures.(8) **(EG 2)**

Alternatives

Most patients are initially given a trial of nonoperative treatment.(1)(3)(9) **(EG 2)**

Length of Stay

Analysis of procedure data for a commercially insured population shows 99% of patients undergoing bunionectomy were discharged the day of or the day after surgery.(21) **(EG 3)** Analysis of procedure data for a Medicare-insured population shows 98% of patients undergoing bunionectomy were discharged the day of or

the day after surgery.(21) (EG 3)

Rationale

Use of this MCG care guideline helps the clinician identify, for a given procedure, which patient-specific factors and clinical conditions are appropriate for that procedure. The evidence-based clinical criteria assist the clinician in the decision to appropriately perform a procedure, evaluating whether the potential benefits of a procedure outweigh the potential risks. For Medicare enrollees, surgical MCG care guidelines also identify which procedures CMS has designated as inpatient only.

Use of these evidence-based clinical criteria to support decision making around the need for a given procedure is of benefit to the patient, as all procedures come with inherent risk that must be balanced by anticipated clinical benefit. Utilizing evidence-based clinical criteria enables a more accurate and patient-specific decision-making process. In addition, the use of evidence-based guidelines can help reduce unwarranted variation in care, such as divergent clinical thresholds to perform a procedure for clinically similar patients that vary across geographic regions, between facilities, and among individual clinicians.

Related CMS Coverage Guidance

This guideline supplements but does not replace, modify, or supersede existing Medicare regulations or applicable National Coverage Determinations (NCDs) or Local Coverage Determinations (LCDs).

Code of Federal Regulations (CFR): 42 CFR 412.3(44); 42 CFR 419.22(45); 42 CFR 422.101(46)

Internet-Only Manual (IOM) Citations: CMS IOM Publication 100-02, Medicare Benefit Policy Manual, Chapter 1 - Inpatient Hospital Services Covered Under Part A(47); CMS IOM Publication 100-02, Medicare Benefit Policy Manual, Chapter 6 - Hospital Services Covered Under Part B(48); CMS IOM Publication 100-02, Medicare Benefit Policy Manual, Chapter 15 - Covered Medical and Other Health Services(49); CMS IOM Publication 100-08, Medicare Program Integrity Manual, Chapter 6, Section 6.5 - Medical Review of Inpatient Hospital Claims for Part A Payment(50)

Medicare Coverage Determinations: Medicare Coverage Database(51)

References

1. Murphy GA. Disorders of the hallux. In: Azar FM, Beaty JH, editors. Campbell's Operative Orthopaedics. 14th ed. Philadelphia, PA: Elsevier; 2021:4041-4153.e7. [Context Link 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13]
2. Meyr AJ, et al. The American College of Foot and Ankle Surgeons® clinical consensus statement: hallux valgus. Journal of Foot and Ankle Surgery 2022 Mar-Apr;61(2):369-383. DOI: 10.1053/j.jfas.2021.08.011. [Context Link 1, 2] View abstract...
3. Hunt KJ. Hallux valgus. In: Haskell A, Coughlin MJ, editors. Coughlin and Mann's Surgery of the Foot and Ankle. 10th ed. Elsevier; 2024:151-274. [Context Link 1, 2, 3, 4, 5, 6]
4. Dias CG, Godoy-Santos AL, Ferrari J, Ferretti M, Lenza M. Surgical interventions for treating hallux valgus and bunions. Cochrane Database of Systematic Reviews 2024, Issue 7. Art. No.: CD013726. DOI: 10.1002/14651858.CD013726.pub2. [Context Link 1] View abstract...
5. Ettinger S, Spindler FT, Savli M, D.A.F. Scientific Committee, Baumbach SF. Correction potential and outcome of various surgical procedures for hallux valgus surgery: a living systematic review and meta-analysis. Archives of Orthopaedic and Trauma Surgery 2024;144(11):4725-4736. DOI: 10.1007/s00402-024-05521-0. [Context Link 1] View abstract...
6. Shi GG, Humayun A, Whalen JL, Kitaoka HB. Management of bunionette deformity. Journal of the American Academy of Orthopedic Surgeons 2018;26(19):e396-e404. DOI: 10.5435/JAAOS-D-17-00345. [Context Link 1, 2, 3, 4] View abstract...
7. Shi GG, Whalen JL, Turner NS, Kitaoka HB. Operative approach to adult hallux valgus deformity: principles and techniques. Journal of the American Academy of Orthopedic Surgeons 2020;28(10):410-418. DOI: 10.5435/JAAOS-D-19-00324. [Context Link 1, 2, 3, 4, 5] View abstract...

8. York PJ, Wydra FB, Hunt KJ. Injuries to the great toe. *Current Reviews in Musculoskeletal Medicine* 2017;10(1):104-112. DOI: 10.1007/s12178-017-9390-y. [Context Link 1, 2] View abstract...
9. Mahan ST, Cidambi EO. Juvenile hallux valgus. *Foot and Ankle Clinics* 2021;26(4):807-828. DOI: 10.1016/j.fcl.2021.07.008. [Context Link 1, 2, 3, 4, 5] View abstract...
10. Seidenstein AH, Torrez TW, Andrews NA, Patch DA, Conklin MJ, Shah A. Pediatric hallux valgus: An overview of history, examination, conservative, and surgical management. *Paediatrics Child Health* 2022;27(2):75-81. DOI: 10.1093/pch/pxab074. [Context Link 1, 2, 3, 4, 5, 6, 7] View abstract...
11. Buddecke DE, Reese ER, Prusa RD. Revision of malaligned lapidus and nonunited lapidus. *Clinics in Podiatric Medicine and Surgery* 2020;37(3):505-520. DOI: 10.1016/j.cpm.2020.03.010. [Context Link 1] View abstract...
12. Migliorini F, Eschweiler J, Tingart M, Maffulli N. Revision surgeries for failed hallux valgus correction: A systematic review. *Surgeon* 2021;19(6):e497-e506. DOI: 10.1016/j.surge.2020.11.010. [Context Link 1] View abstract...
13. Trnka HJ. Managing complications of foot and ankle surgery: hallux valgus. *Foot and Ankle Clinics* 2022;27(2):271-285. DOI: 10.1016/j.fcl.2021.11.015. [Context Link 1] View abstract...
14. Barnes D, et al. From etiology to intervention: a holistic review of bunion pathophysiology and care. *Advances in Orthopedics* 2024;2024:9910410. DOI: 10.1155/2024/9910410. [Context Link 1, 2, 3] View abstract...
15. Colo G, Leighab M, Surace MF, Fusini F. The efficacy of shoes modification and orthotics in hallux valgus deformity: a comprehensive review of literature. *Musculoskeletal Surgery* 2024;Online. DOI: 10.1007/s12306-024-00839-9. [Context Link 1, 2] View abstract...
16. Krzesniak H, Truszczynska-Baszak A. Toe separators as a therapeutic tool in physiotherapy-a systematic review. *Journal of Clinical Medicine* 2024;13(24):DOI: 10.3390/jcm13247771. [Context Link 1] View abstract...
17. AlZeedi M, Park JP, Marwan Y, Abu-Dalu KM, Hamdy R, Janelle C. Growth modulation for the treatment of juvenile hallux valgus: a systematic review of literature. *Strategies in Trauma and Limb Reconstruction* 2023;18(1):51-55. DOI: 10.5005/jp-journals-10080-1579. [Context Link 1] View abstract...
18. Artioli E, et al. First metatarsal hemiepiphyodesis for the treatment of juvenile hallux valgus: a systematic review. *Journal of Pediatric Orthopedics* 2023;43(9):584-589. DOI: 10.1097/BPO.0000000000002485. [Context Link 1] View abstract...
19. Lewis TL, Patel K, Shepherd KL, MacInnes P, Ray R, Kokkinakis M. Hallux valgus surgery in children with cerebral palsy: A systematic review. *Foot and Ankle Surgery* 2022;28(4):476-482. DOI: 10.1016/j.fas.2021.12.009. [Context Link 1] View abstract...
20. Attia AK, Heier KA. First metatarsophalangeal arthrodesis for the failed hallux. *Foot and Ankle Clinics* 2022;27(4):723-744. DOI: 10.1016/j.fcl.2022.07.001. [Context Link 1, 2] View abstract...
21. Proprietary health insurance data sources (2023-2024); and Medicare 5% Standard Analytical File (2022-2023). [Context Link 1, 2, 3]
22. Premier PINC AI™ Healthcare Database (PHD), 01/01/2023-12/31/2024. Premier, Inc. [Context Link 1]
23. Hudson T. The role of social determinates of health in discharge practices. *Nursing Clinics of North America* 2021;56(3):369-378. DOI: 10.1016/j.cnur.2021.04.004. [Context Link 1] View abstract...
24. Chovanec KA, Arsene C, Beck A, Zachrich K, Liedel B, Wolff-Elliott J. Association of discharge disposition with outcomes. *Population Health Management* 2021;24(1):116-121. DOI: 10.1089/pop.2019.0176. [Context Link 1] View abstract...
25. Centers for Medicare and Medicaid Services. "Condition of participation: Discharge planning." 42 CFR Pt. 482.43 Washington, DC 2025 Jan 01 [accessed 2025 Jul 22] Accessed at: <https://www.ecfr.gov/>. [Context Link 1, 2, 3]
26. Management of patients with musculoskeletal disorders. In: Hinkle JL, Cheever KH, Overbaugh KJ, editors. *Brunner & Suddarth's Textbook of Medical-Surgical Nursing*. 15th ed. Wolters Kluwer; 2022:1113-1151. [Context Link 1]
27. Orthopedic surgery. In: Hornacky A, Phillips N, editors. *Berry & Kohn's Operating Room Technique*. 15th ed. Elsevier; 2025:761-789. [Context Link 1, 2]
28. Filson R. Musculoskeletal trauma and orthopedic surgery. In: Harding MM, Kwong J, Hagler D, Reinisch C, editors. *Lewis's Medical-Surgical Nursing: Assessment and Management of Clinical Problems*. 12th ed. St. Louis, MO: Mosby; 2023:1638-1672. [Context Link 1, 2]
29. Skin integrity and wound care. In: Lynn P, editor. *Taylor's Clinical Nursing Skills*. 6th ed. Wolters Kluwer; 2023:452-540. [Context Link 1, 2]

30. Targeted care planning and care transitions. In: Perez R, editor. CMSA's Integrated Case Management. 2nd ed. Springer Publishing Company LLC; 2023:178-189. [Context Link 1, 2]
31. Elements of Excellence in Transitions of Care (TOC). TOC Checklist [Internet] National Transitions of Care Coalition. 2006 Accessed at: <https://www.ntocc.org>. [accessed 2025 Sep 16] [Context Link 1]
32. Medical-surgical nursing. In: Hinkle JL, Cheever KH, Overbaugh KJ, editors. Brunner & Suddarth's Textbook of Medical-Surgical Nursing. 15th ed. Wolters Kluwer; 2022:33-55. [Context Link 1]
33. Transition of care. In: Gillingham DC, editor. Foundations of Case Management. Blue Bayou Press; 2021:137-144. [Context Link 1]
34. Becker C, et al. Interventions to improve communication at hospital discharge and rates of readmission: a systematic review and meta-analysis. JAMA Network Open 2021;4(8):e2119346. DOI: 10.1001/jamanetworkopen.2021.19346. [Context Link 1] View abstract...
35. The nursing process in drug therapy and patient safety. In: Tucker RG, editor. Karch's Focus on Nursing Pharmacology. 9th ed. Wolters Kluwer; 2023:47-57. [Context Link 1]
36. Adverse drug reactions and medication errors. In: Burchum JR, Rosenthal LD, editors. Lehne's Pharmacology for Nursing Care. 12th ed. Elsevier; 2025:64-75. [Context Link 1]
37. Safe medication preparation. In: Perry AG, Potter PA, Ostendorf WR, Laplante N, editors. Clinical Nursing Skills and Techniques. 11th ed. Elsevier; 2025:586-607. [Context Link 1]
38. The case manager's roles, functions, and activities. In: Gillingham DC, editor. Foundations of Case Management. Blue Bayou Press; 2021:9-14. [Context Link 1]
39. Ioli JP, Sandberg-Cook J. Ankle and foot pain. In: Buttaro TM, Polgar-Bailey P, Sandberg-Cook J, Dick KL, Montgomery JB, editors. Primary Care: Interprofessional Collaborative Practice. 7th ed. Elsevier; 2025:1022-1030. [Context Link 1, 2]
40. Anderson TS, Ayanian JZ, Herzig SJ, Souza J, Landon BE. Gaps in primary care follow-up after hospital discharge among Medicare beneficiaries. Journal of the American Geriatrics Society 2025;DOI: 10.1111/jgs.19496. [Context Link 1] View abstract...
41. Bragg-Underwood T. Family health. In: Nies MA, editor. Community/Public Health Nursing. 8th ed. Elsevier; 2024:404-427. [Context Link 1]
42. Harding MM. Substance use disorders in acute care. In: Harding MM, Kwong J, Hagler D, Reinisch C, editors. Lewis's Medical-Surgical Nursing: Assessment and Management of Clinical Problems. 12th ed. St. Louis, MO: Mosby; 2023:162-178. [Context Link 1]
43. Management of patients with dermatologic disorders. In: Hinkle JL, Cheever KH, Overbaugh KJ, editors. Brunner & Suddarth's Textbook of Medical-Surgical Nursing. 15th ed. Wolters Kluwer; 2022:1815-1864. [Context Link 1]
44. Centers for Medicare and Medicaid Services. "Admissions." 42 CFR 412.3 Washington, DC 2025 Jul [accessed 2025 Jul 22] Accessed at: <https://www.ecfr.gov/current/title-42/chapter-IV/subchapter-B/part-412>. [Context Link 1]
45. Centers for Medicare and Medicaid Services. "Hospital services excluded from payment under the hospital outpatient prospective payment system." 42 CFR 419.22 Washington, DC 2023 Jul [accessed 2025 Jul 22] Accessed at: <https://www.ecfr.gov/>. [Context Link 1]
46. Centers for Medicare and Medicaid Services. "Requirements relating to basic benefits." 42 CFR 422.101 Washington, DC 2025 Jun 03 [accessed 2025 Jul 23] Accessed at: <https://www.ecfr.gov/>. [Context Link 1]
47. Centers for Medicare and Medicaid Services. Medicare Benefit Policy Manual. Chapter 1 - Inpatient Hospital Services Covered Under Part A Rev. 10892 [Internet] Centers for Medicare and Medicaid Services. 2021 Aug Accessed at: <https://www.cms.gov/manuals/>. [accessed 2025 Sep 09] [Context Link 1]
48. Centers for Medicare and Medicaid Services. Medicare Benefit Policy Manual. Chapter 6 - Hospital Services Covered Under Part B Rev. 12425 [Internet] Centers for Medicare and Medicaid Services. 2023 Dec Accessed at: <http://www.cms.gov/manuals/>. [accessed 2025 Sep 09] [Context Link 1]
49. Centers for Medicare and Medicaid Services. Medicare Benefit Policy Manual. Chapter 15 - Covered Medical and Other Health Services Rev. 13108 [Internet] Centers for Medicare and Medicaid Services. 2025 Apr 11 Accessed at: <https://www.cms.gov/manuals/>. [accessed 2025 Sep 09] [Context Link 1]
50. Centers for Medicare and Medicaid Services. Medicare Program Integrity Manual. Chapter 6, Section 6.5 - Medical Review of Inpatient Hospital Claims for Part A Payment Rev. 10365 [Internet] Centers for Medicare and Medicaid Services. 2020 Oct 02 Accessed at: <https://www.cms.gov/medicare/regulations-guidance/manuals/internet-only-manuals-ioms>. [accessed 2025 Sep 09] [Context Link 1]

51. Medicare Coverage Database. [Internet] Centers for Medicare and Medicaid Services. Accessed at: <https://www.cms.gov/medicare-coverage-database/search.aspx>? Updated 2025 [accessed 2025 Oct 23] [Context Link 1]

Footnotes

[A] See Ambulatory Surgery Discharge and Complications: Common Complications and Conditions  ISC for further information. [A in Context Link 1]

[B] Expert consensus varies as to whether or not non-weight-bearing immobilization following bunionectomy is beneficial.(2) [B in Context Link 1]

[C] Some patients may have their hydration needs met via alternative means (eg, percutaneous endoscopic gastrostomy tube). [C in Context Link 1]

[D] Discharge instructions should be given in the patient's and caregiver's native language using trained language interpreters whenever possible.(23) [D in Context Link 1]

Definitions

Hemodynamic stability

- Hemodynamic stability, as indicated by **1 or more** of the following:
 - Hemodynamic abnormalities at baseline or acceptable for next level of care
 - Patient hemodynamically stable, as indicated by **ALL** of the following(1)(2):
 - Tachycardia absent
 - Hypotension absent
 - No evidence of inadequate perfusion (eg, no myocardial ischemia)
 - No other hemodynamic abnormalities (eg, no Orthostatic hypotension)

References

1. Schrager DL. Approach to the patient with abnormal vital signs. In: Goldman L, Cooney KA, editors. Goldman-Cecil Medicine. 27th ed. Elsevier; 2024:32-35.
2. Ramgopal S, Sepanski RJ, Martin-Gill C. Empirically derived age-based vital signs for children in the out-of-hospital setting. Annals of Emergency Medicine 2023;81(4):402-412. DOI: 10.1016/j.annemergmed.2022.09.019.

Hypotension absent

- Hypotension absent,[A] as indicated by **1 or more** of the following:
 - Hypotension absent in adult patient, as indicated by **1 or more** of the following:
 - Systolic blood pressure greater than or equal to 90 mm Hg[A](1)
 - Mean arterial pressure[B] greater than or equal to 70 mm Hg  MAP Calculator[A](1)(2)
 - Blood pressure at patient's baseline (eg, healthy adult with low systolic blood pressure), at intentional therapeutic goal (eg, patient with heart failure), or acceptable for next level of care (eg, blood pressure stable and no significant signs or symptoms due to low blood pressure)
 - Hypotension absent in pediatric patient, as indicated by **1 or more** of the following:
 - Systolic blood pressure greater than or equal to 110 mm Hg in child 13 to 17 years of age[A](3)
 - Systolic blood pressure greater than or equal to 100 mm Hg in child 6 to 12 years of age[A](3)
 - Systolic blood pressure greater than or equal to 95 mm Hg in child 3 to 5 years of age[A](3)

- Systolic blood pressure greater than or equal to 90 mm Hg in child 1 or 2 years of age^{[A](3)}
- Systolic blood pressure greater than or equal to 80 mm Hg in infant 6 to 11 months of age^{[A](3)}
- Systolic blood pressure greater than or equal to 70 mm Hg in infant 3 to 5 months of age^{[A](3)}
- Systolic blood pressure greater than or equal to 65 mm Hg in infant 1 or 2 months of age^{[A](3)}
- Blood pressure at patient's baseline (eg, healthy child with low systolic blood pressure), at intentional therapeutic goal, or acceptable for next level of care (eg, blood pressure stable and no significant signs or symptoms due to low blood pressure)

References

1. Schriger DL. Approach to the patient with abnormal vital signs. In: Goldman L, Cooney KA, editors. Goldman-Cecil Medicine. 27th ed. Elsevier; 2024:32-35.
2. Puskasich MA, Jones AE. Shock. In: Walls RM, editor. Rosen's Emergency Medicine: Concepts and Clinical Practice. 10th ed. Elsevier; 2023:34-41.e1.
3. Anderson CC, Kapoor S, Mark TE. Pediatric parameters and equipment. In: Anderson CC, Kapoor S, Mark TE, editors. The Harriet Lane Handbook: A Manual for Pediatric House Officers. 23rd ed. Elsevier; 2024:i-iii.

Footnotes

- A. Criteria based upon clinician-acquired numeric values (eg, vital signs, oxygen saturation) should be used if they are accurate reflections of the patient's condition. Transitory findings (eg, abnormal only upon initial emergency department intake or only one time out of multiple readings) that rapidly improve with no or minimal treatment usually do not reflect disease severity or risk for deterioration. This does not imply that an initial or one-time reading cannot ever be applicable. The goal is to separate erroneous or incidental findings from those that truly represent the patient's clinical picture.
- B. The mean arterial pressure (MAP) takes into account both SBP and DBP readings.

Multimodal analgesia

- Multimodal analgesia involves the utilization of 2 or more analgesic agents with different mechanisms of action in order to provide additive or synergistic pain control, while minimizing side effects and reliance on opioids.⁽¹⁾⁽²⁾⁽³⁾

References

1. D'Souza RS, Johnson RL. Regional and multimodal treatments of perioperative pain. In: Benzon H, editor. Practical Management of Pain. 6th ed. Philadelphia, PA 19103-2899: Elsevier; 2023:355-373.e8.
2. George S, Johns M. Review of nonopioid multimodal analgesia for surgical and trauma patients. American Journal of Health-System Pharmacy 2020;77(24):2052-2063. DOI: 10.1093/ajhp/zxaa301.
3. Finnerup NB. Nonnarcotic methods of pain management. New England Journal of Medicine 2019;380(25):2440-2448. DOI: 10.1056/NEJMra1807061.

Orthostatic hypotension

- Orthostatic hypotension,^{[A][B]} as indicated by **1 or more** of the following⁽¹⁾⁽²⁾⁽³⁾:
 - Fall in SBP of 20 mm Hg or more 1 to 3 minutes after patient sits or stands from recumbent position
 - Fall in DBP of 10 mm Hg or more 1 to 3 minutes after patient sits or stands from recumbent position

References

1. Shibao C, Lipsitz LA, Biaggioni I, American Society of Hypertension Writing Group. Evaluation and treatment of orthostatic hypotension. Journal of the American Society of Hypertension 2013 Jul-Aug;7(4):317-324. DOI: 10.1016/j.jash.2013.04.006.

2. Dalal AS, Van Hare GF. Syncope. In: Kliegman RM, et al., editors. Nelson Textbook of Pediatrics. 22nd ed. Elsevier; 2025:592-599.
3. Fang JC, O'Gara PT. History and physical examination: an evidence-based approach. In: Libby P, Bonow RO, Mann DL, Tomaselli GF, Bhatt DL, Solomon SD, editors. Braunwald's Heart Disease: A Textbook of Cardiovascular Medicine. 12th ed. Elsevier; 2022:123-140.

Footnotes

- A. Concomitant measurements of the heart rate are important to measure to help diagnose subtypes of orthostatic hypotension (eg, the lack of a compensatory increase in heart rate is typical of autonomic failure and an exaggerated tachycardia may be reflective of volume depletion). However, the heart rate is not a component of the definition of orthostatic hypotension, which relies upon blood pressure alone.(1)(2)(3)
- B. Criteria based upon clinician acquired numeric values (eg, vital signs, oxygen saturation) should be used if they are accurate reflections of the patient's condition. Transitory findings (eg, abnormal only upon initial emergency department intake or only one time out of multiple readings) that rapidly improve with no or minimal treatment usually do not reflect disease severity or risk for deterioration. This does not imply that an initial or one-time reading cannot ever be applicable. The goal is to separate erroneous or incidental findings from those that truly represent the patient's clinical picture.

Social Determinants of Health Assessment

- Risk of poor health outcomes may be increased by the presence of **1 or more** of the following social determinants of health(1)(2)(3):
 - Housing insecurity, as indicated by **1 or more** of the following:
 - Individual or caregiver's current living situation is **1 or more** of the following(4):
 - Does not have own housing (eg, staying in a hotel, shelter, or with others)
 - Has own housing (eg, house, apartment), but at risk of losing it in the future (ie, behind on rent or mortgage)
 - Has own housing (eg, house, apartment), but has lived in 3 or more places in past year
 - Current housing has **1 or more** of the following:
 - Electrical appliances (eg, stove, refrigerator) not working or unavailable
 - Insufficient heating or cooling
 - Insufficient ventilation
 - Lead paint or pipes
 - Mold
 - Pests (eg, bugs) or rodents
 - Smoke detectors not working or unavailable
 - Food insecurity, as indicated by **1 or more** of the following(5):
 - In the past year, individual or caregiver ran out of food and did not have money to buy more food.
 - In the past year, individual or caregiver worried that they would run out of food before they received money to buy more food.
 - Insufficient transportation, as indicated by **1 or more** of the following(6):
 - In the past year, individual or caregiver missed medical appointments or could not get medications due to lack of transportation.
 - In the past year, individual or caregiver missed nonmedical activities, work, or could not get things needed for daily living due to lack of transportation.
 - Insufficient utilities, as indicated by **1 or more** of the following(7):
 - Utilities (eg, electricity, water, gas, or oil) are currently shut off or unavailable.
 - In the past year, electric, water, gas, or oil company threatened to shut off services.
 - Personal safety risk, as indicated by **2 or more** of the following(5):
 - Individual is sometimes or frequently physically hurt by another person (including family member).
 - Individual is sometimes or frequently insulted or talked down to by another person (including family member).

- Individual is sometimes or frequently threatened with physical harm by another person (including family member).
- Individual is sometimes or frequently screamed or cursed at by another person (including family member).
- Insufficient dependent care, as indicated by **1 or more** of the following:
 - In the past year, individual or caregiver was unable to work due to lack of dependent care.
 - In the past year, individual or caregiver was unable to work more (additional) hours due to lack of dependent care.
 - In the past year, individual or caregiver missed medical appointments or could not get medications due to lack of dependent care.
 - In the past year, individual or caregiver missed nonmedical activities (eg, school, church, social activity) due to lack of dependent care.
- Depression risk, as indicated by **ALL** of the following(8):
 - In the past 2 weeks, individual had little interest or pleasure in normal activities on at least several days.
 - In the past 2 weeks, individual felt down, depressed, or hopeless on at least several days.

References

1. Scanlon A, Reinisch C. Social determinants of health. In: Harding MM, Kwong J, Hagler D, Reinisch C, editors. Lewis's Medical-Surgical Nursing: Assessment and Management of Clinical Problems. 12th ed. St. Louis, MO: Mosby; 2023:18-20.
2. Billioux A, Verlander K, Anthony S, Alley D. Standardized Screening for Health-Related Social Needs in Clinical Settings: the Accountable Health Communities Screening Tool. [Internet] National Academy of Sciences. 2017 May Accessed at: <https://nam.edu/>. [accessed 2025 Sep 12]
3. Addressing social determinants of health. In: Perez R, editor. CMSA's Integrated Case Management. 2nd ed. Springer Publishing Company LLC; 2023:62-75.
4. Sandel M, et al. Unstable housing and caregiver and child health in renter families. Pediatrics 2018;14(2):e20172199. DOI: 10.1542/peds.2017-2199.
5. Children's HealthWatch Survey. Screening Instrument [Internet] Children's HealthWatch. 2024 Feb 13 Accessed at: <https://childrenshealthwatch.org/>. [accessed 2025 Mar 31]
6. PRAPARE®: Protocol for Responding to and Assessing Patient Assets, Risks, and Experiences Screening Tool. 2.0 [Internet] Association of Asian Pacific Community Health Organizations (AAPCHO) and National Association of Community Health Centers (NACHC). 2025 Accessed at: <https://prapare.org/the-prapare-screening-tool/>. [accessed 2025 Sep 08]
7. Cook JT, et al. A brief indicator of household energy security: associations with food security, child health, and child development in US infants and toddlers. Pediatrics 2008;122(4):e867-75. DOI: 10.1542/peds.2008-0286.
8. Kroenke K, Spitzer RL, Williams JB. The Patient Health Questionnaire-2: validity of a two-item depression screener. Medical Care 2003;41(11):1284-92. DOI: 10.1097/01.MLR.0000093487.78664.3C.

Tachycardia absent

- Tachycardia[A][B] absent, as indicated by **1 or more** of the following:
 - Heart rate less than or equal to 100 beats per minute in adult[A][B](1)
 - Heart rate less than or equal to 85 beats per minute in child 13 to 17 years of age[A][B](2)
 - Heart rate less than or equal to 95 beats per minute in child 6 to 12 years of age[A][B](2)
 - Heart rate less than or equal to 110 beats per minute in child 1 to 5 years of age[A][B](2)
 - Heart rate less than or equal to 120 beats per minute in infant 3 to 11 months of age[A][B](2)
 - Heart rate less than or equal to 150 beats per minute in infant 1 or 2 months of age[A][B](2)

References

1. Schriger DL. Approach to the patient with abnormal vital signs. In: Goldman L, Cooney KA, editors. Goldman-Cecil Medicine. 27th ed. Elsevier; 2024:32-35.
2. Anderson CC, Kapoor S, Mark TE. Pediatric parameters and equipment. In: Anderson CC, Kapoor S, Mark TE, editors. The Harriet Lane Handbook: A Manual for Pediatric House Officers. 23rd ed. Elsevier; 2024:i-iii.

Footnotes

- A. Criteria based upon clinician acquired numeric values (eg, vital signs, oxygen saturation) should be used if they are accurate reflections of the patient's condition. Transitory findings (eg, abnormal only upon initial emergency department intake or only one time out of multiple readings) that rapidly improve with no or minimal treatment usually do not reflect disease severity or risk for deterioration. This does not imply that an initial or one-time reading cannot ever be applicable. The goal is to separate erroneous or incidental findings from those that truly represent the patient's clinical picture.
- B. Interpretation of heart rate requires clinical judgment and consideration of several patient-specific factors, such as the patient's baseline heart rate, medications, and clinical impact. For example, an elderly patient on a beta-blocker medication with a baseline resting heart rate of 60 beats per minute may be clinically tachycardic at a heart rate of 94 beats per minute. Likewise, a patient who is upset, in pain, or nervous in the emergency department with a heart rate of 106 beats per minute may meet the technical definition of tachycardia, but this tachycardia (absent associated findings such as chest pain or hypotension) may not be clinically important. The numeric values included in this definition are provided to allow for consistency in terms of a technical definition of the term tachycardia. Whether a heart rate above or below the technical threshold is clinically meaningful is a matter of persistence, context, and clinical judgment.

Codes

ICD-10 Diagnosis: M20.10, M20.11, M20.12, M21.611, M21.612, M21.619, M21.621, M21.622, M21.629

ICD-10 Procedure: 0MQS0ZZ, 0MQT0ZZ, 0Q8N0ZZ, 0Q8P0ZZ, 0QBN0ZZ, 0QBP0ZZ, 0QQN0ZZ, 0QQP0ZZ, 0QQQ0ZZ, 0QQR0ZZ, 0QSN042, 0QSN04Z, 0QSN052, 0QSN342, 0QSN352, 0QSP042, 0QSP04Z, 0QSP052, 0QSP342, 0QSP352, 0QSQ04Z, 0QSR04Z, 0SRP0JZ, 0SRQ0JZ, 0STP0ZZ, 0STQ0ZZ [Hide]

CPT®: 28292, 28295, 28296, 28297, 28298, 28299

CPT copyright 2025 American Medical Association. All rights reserved.

MCG Health
Inpatient & Surgical Care 30th Edition
Copyright © 2026 MCG Health, LLC
All Rights Reserved

Last Update: 1/25/2026 6:19:01 AM
Build Number: 30.0.2026012500524.025256